



# COMMON CAMAS

## *Camassia quamash* (Pursh) Greene ssp. *breviflora* Gould

Plant Symbol = CAQUB2

*Common Names:* small camas, blue camas

*Scientific Names:* *Camassia quamash* (Pursh) Greene var. *breviflora* (Gould) C.L. Hitchc.

**Warning:** Death camas (*Zigadenus venenosus*) can be confused with edible camas bulbs and is toxic. Be sure of your identification of camas bulbs before eating them!

### Description

*General:* Lily Family (Liliaceae). Common camas is a stout, robust plant 12-28 inches (30-70 cm) tall with a dense inflorescence. It is a perennial herb that grows from an edible bulb. The leaves are long and narrow, grass-like, and emerge from the base. Common camas flowers are light to deep blue; more than three flowers in an inflorescence may be open at one time. Camas flowers have six tepals, six stamens, and three stigmas. The inflorescence is a spike-like cluster borne on a leafless stem that is held above the leaves. Common camas is distinguished from great camas (*Camassia leichtlinii* ssp. *suksdorfii*) by the following: the flowers are slightly irregular, with the lowest tepal curving outward away from the stem; the anthers are bright yellow; the plant is relatively short and stout, with shorter flower stalks and smaller bulbs; and there is no waxy powder on the leaves. Common camas blooms from April through June. The fruits are barrel-shaped to three-angled capsules, splitting into three parts to release many black, angled seeds.

*Distribution:* *Camassia quamash* ssp. *breviflora* is found in the West from British Columbia through Washington, Oregon, Nevada, and California at elevations below 8000 ft (2500 m). The southern limit of its range in California is the high Sierra Nevada Mountains and Modoc Plateau. In British Columbia, common camas is found in moist meadows, rocky outcrops, bluffs, and islands in southwestern British Columbia, mainly on southwestern Vancouver Island and the Gulf Islands. For current distribution, please consult the Plant Profile page for this species on the PLANTS Web site.

*Habitat:* Common camas grows in wet meadows, wet prairies, swales, depressions, annual floodplains, moist hillsides, and streamside areas. Camas habitat is often ephemeral, drying out by late spring.

### Adaptation

Common camas is adapted to seasonally wet soils in full sun to partial shade. Bulb depth appears limited by shallow water tables, anoxic conditions, or restrictive layers.

### Uses

*Ornamental:* Horticulturally, this plant is used for cut flowers, beds, borders, ground cover, rock gardens, and prairie restoration.

*Wildlife:* Elk, deer, and moose reportedly graze camas early in the spring (Craighead et al., 1963). Gophers eat camas and move the bulbs to other areas where they sprout and grow the next year (Watson, 1988). Indian women in Oregon's Umpqua Valley robbed camas bulbs from gopher caches (Piper, 1916). Herbivorous insects also eat camas leaves. Camas attracts a wide variety of pollinators including European honeybees, bumble bees, mason bees, hover flies, beetles, and lady beetles (Bartow, 2015; Pendergrass et al., 2008).



Photo by Kathy Pendergrass, Oregon NRCS, 2007.

## Ethnobotany

Camas was and continues to be one of the most important "root" foods of western North American indigenous peoples, from southwestern British Columbia to Montana, and south to California (Kuhnlein and Turner, 1991). The part of the plant that was relished is actually a bulb. Camas was used by Northwest Coast peoples, the Coast Salish of Vancouver Island, western Washington groups, Squamish, Sechelt, Comox, and Kwak-waka'wakw of the British Columbia coast. Camas was considered to be one of the most important bulbs to local California natives. The Maidu particularly valued great camas.

Except for choice varieties of dried salmon, no other food item was more widely traded (Gunther, 1973). People traveled great distances to harvest the bulbs and there is some suggestion that plants were dispersed beyond their range by transplanting (Turner and Efrat, 1982; Turner et al., 1983). To the Nez Perce people, camas is still the most important root in trade, and trading is traditionally impossible without camas bulbs (Harbinger, 1964). Dried camas is the most expensive form of camas, with baked and then raw camas being less expensive. At marriage trades, the girl's family gives roots in corn husk bags. At funeral trades, the widow gives camas roots to friends and relatives. The Nez Perce traded camas roots with the Warm Springs, Umatilla, Cayuse, Walla Walla, Nespelem, Yakama, Crows, and Flatheads.

The bulbs were usually dug after flowering, in summer, although some peoples dug them in spring. Harvesting the bulbs traditionally took weeks or months among the Nez Perce. Each family group "owned" its own camping spot and harvesting spot. These were passed down in families from generation to generation. Turf was lifted out systematically in small sections and then replaced after only larger bulbs had been removed. The bulbs were dug with a pointed digging stick. Annual controlled burning was used to maintain an open prairie-like habitat for optimum camas production. Areas were harvested only every few years.

Traditionally, camas bulbs were almost always pit-cooked. Within the past 100 years, camas bulbs have also been cooked by stovetop methods (Turner and Kuhnlein, 1983). The bulbs are allowed to cook for 24-36 hours when pit-cooked (Turner and Bell, 1971). It is probable that lengthy cooking is necessary for maximum conversion of the inulin in *Camassia* to fructose. The sweetness of cooked camas gave it utility as a sweetener and enhancer of other foods. Before sugar, European traders introduced molasses, and honey. Sweetening agents were in short supply among native peoples, and camas was highly valued in this capacity. Sometimes other foods, such as the rhizomes of springbank clover (*Trifolium wormskioldii*) and the roots of Pacific silverweed (*Potentilla anserina* ssp. *pacifica*) were cooked with the camas bulbs. The Kalapuyan of the Willamette Valley in Oregon used to flavor camas with tarweed (*Madia elegans*). Bulbs don't keep well fresh. They were cooked or sun-dried and stored for later use. Sometimes camas bulbs were pressed flat and made into camas cakes the size of biscuits before being dried (Turner et al., 1983). Dried bulbs were re-constituted by soaking in water, usually overnight.

Many of the traditional camas gathering sites, such as Weippe Prairie and Camas Prairie in Idaho and the Willamette Valley in Oregon, have been converted to agriculture. The average size of a camas patch needed to feed a five person family was 6.7 acres (Thoms, 1989). Camas roots are hard to find now. Restoration of camas prairies and access to camas bulbs are priorities of many Indian people. At one time, "When camas was in bloom in wet meadows, the flowers grow so thickly that they look like a blue lake" (Murphey, 1959).

Camas stalks and leaves were used for making mattresses. It was sometimes used in place of grass when baking camas in pits. Camas is used by the Nez Perce as a cough medicine. It is boiled, and the juice is strained and mixed with honey.

*Traditional Resource Management* (TRM) was often intensive, to the point of being considered "semi-agricultural" by some. According to Dr. Nancy Turner, TRM included the following:

- Ownership, demarcation, and inheritance of beds or patches,
- Clearing of rock, brush, and weedy vegetation,
- Harvesting bulbs after seeds were produced, during specific times of the year,
- Periodic field burning in summer after digging,



*Common camas and rosy seablush (Plectritis congesta) blooming in the Camassia Natural Area in western Oregon. Photo by Kathy Pendergrass, Oregon NRCS, 2007.*

- In some cases, sod removal then bulb removal followed by sod replacement,
- Digging or “cultivation” to keep the soil loose,
- “Selective breeding” by transplanting “better” bulbs to the beds,
- Sustainable harvest techniques, including partial, selective harvests and incidental or planned promotion of camas colonization and reproduction, and
- Death camas bulbs (*Zigadenus venenosus*) were removed, so they wouldn’t accidentally be mistaken for the edible camas bulbs.

### **Status**

*Wetland Indicator:* Common camas is classified as a facultative wetland plant (FACW), meaning it usually occurs in wetlands, but also may occur in non-wetlands. Please consult the PLANTS Web site (<http://plants.usda.gov/>) and your state’s Department of Natural Resources for this plant’s current status (e.g., threatened or endangered species, state noxious status, and wetland indicator values).

### **Planting Guidelines**

Common camas can be propagated from seeds or bulbs. Common camas generally prefers full sun to partial shade, with bulb depth ranging from 2-8 inches (most commonly 4-6 inches deep). The bulbs of common camas can be substantially smaller in size and occur at more shallow soil depths than great camas. The occasional occurrence of a large, thick root beneath a bulb may aid in re-locating or re-establishing it at a greater depth. Plants require irrigation or moist soil conditions to become established, and camas can be difficult to establish in California.

Common camas is readily established by transplanting wild or commercially grown bulbs. Wild harvests should be restricted to salvage sites with appropriate approvals or permits. Due to loss of wetland habitat throughout the United States, harvesting plants from the wild is rarely appropriate or legal except under salvage situations. Use of bulbs or seeds from local nurseries or greenhouses is strongly recommended.

Given that camas commonly occupies sites high in silt and clay that dry out in summer, windows for digging are often narrow. The best time to excavate bulbs is from early summer through mid-fall. Store the bulbs in a dry, dark, cool, well ventilated place in a potting medium such as dry peat moss, similar to recommendations for fall planted/spring flowering bulbs (such as daffodils and tulips).

Plant camas outdoors in the fall or early winter, when soils are moist enough to dig and prevailing soil temperatures are cool. This is generally below 60°F. Fall planting allows for better root development and fulfillment of any chilling requirement for flowering (De Hertogh et. al. 1993). The larger the bulb, the greater the planting depth can be. As a general rule, bulbs can be planted three times as deep of the diameter of the bulb. Keep the camas bed damp until it gets warm. Once plants senesce after flowering, stop watering so seeds form and bulbs cure. Suggested spacing for flower beds and naturalized landscapes vary from 3-4 inches apart (8-10 per sq. ft.) to 6-8 inches apart.

For prairie restoration, seed common camas in the fall at a single species rate of about 22 pounds per acre (50 seeds per square foot), or if included as part of a pollinator or prairie seed mix, at a rate of 1 to 5 lb/ac (or about 5-10% of the mix, calculated on a seeds-per-square-foot basis).

### **Management**

Consistent soil moisture is required every spring, but the soil can be allowed to dry out soon after the pods mature or the leaves senesce (dry up and turn brown). Moderate soil nutrient levels are beneficial. In natural settings, minor soil disturbance (loosening, surface scarification) adjacent to existing specimens may enhance natural regeneration by seed; seedling recruitment is highest in areas that are bare when the seeds germinate in February. Late summer field burning (where and when permitted) may improve stand vigor, reduce competition from brush and certain weeds, and aid in regeneration. For optimal bulb development, avoid mowing or grazing more than lightly, if at all, even during foliar senescence. Individual plants may live 15 to 20 years or more.

### **Pests and Potential Problems**

Camas is favored as forage by deer, so fencing or repellents may be useful, particularly during the first growing season. Bulb predation by pocket gophers can be a substantial threat to stands if left unchecked. For ornamental use, it may be necessary to bury bulbs with a protective wire mesh to prevent herbivory by voles and pocket gophers. The mesh needs to be coarse enough to allow shoots to grow through (De Hertogh et al., 1993). Voles (field mice) forage on young leaves in early spring and also consume bulbs. It may also be necessary to protect young seedlings from slugs the first year.

### **Environmental Concerns**

There are no known environmental concerns related to common camas.



## Seeds and Plant Production

Common camas propagates easily from seed. It can be collected as soon as the pods mature (turn light brown) or split open to reveal the mature black seeds. Pods ripen from late May to July. Dry seeds can be stored in a cool, dry place prior to planting. Camas seed is dormant and typically requires 60-100 days of cold moist stratification (moist chilling at 34-40°F) for maximum germination (Bartow, 2015; Emery, 1988; Guerrant and Raven, 1995; Deno, 1993; Northway pers. comm. 1998; Thoms, 1989).

To establish seed production fields, sow seed in the fall at a rate of 6 to 10 pounds per acre in 6- to 14-inch wide rows at a depth of ½ to ¼ inch (Bartow, 2015). Fall sown fields can be sprayed with a non-specific herbicide to eliminate the first flush of weeds, since seedlings do not typically germinate until late winter. One-leafed, grass-like seedlings will emerge in February or March under suitable conditions. Seedlings require moisture through the spring growing period to survive. Warm temperatures during seedling development can be lethal.

If started from seed, common camas will not flower and set seed until the third or fourth growing season. Once the bulbs are large enough to flower, seed yields are likely to increase substantially for several years thereafter, eventually plateauing and then starting to decline as bulbs become more crowded. As stands age, an abundance of foliage with few flowers is an indication that the bulbs have become over-crowded; on a small scale, they can be lifted, separated, and replanted at the proper spacing in the fall.

Seed can be harvested by conventional direct combining followed by seed drying (Bartow, 2015). Harvest should occur when the large majority of capsules have turned from green to tan and only a small percentage have split open to reveal the black seeds. If substantial healthy green foliage is still present at harvest time, plants should be cut high to leave as much of the leaves intact to benefit further bulb development. Optionally, mature stalks and seed heads can be windrowed onto tarps and then dried for a week or two. The dry capsules should split and release the seed, thus negating the need for threshing, but some shaking is required.

Common camas has approximately 100,000 seeds per pound.

## Cultivars, Improved, and Selected Materials (and area of origin)

Bulbs and/or seed of common camas are readily available through native plant nurseries within its range. Cultivars of common camas are available from the flower bulb industry. *Camassia quamash* 'Orion' is deep blue, while those of 'San Juan Form' are an even deeper, more vibrant blue. There is also a white form (Benzel, 1995). Cultivars should be selected based on the local climate, resistance to local pests, and intended use. Consult with your local land grant university, local extension or local USDA NRCS office for recommendations on adapted cultivars for use in your area.



White form of common camas blooming in Camassia Natural Area in western Oregon. Photo by Kathy Pendergrass, Oregon NRCS, 2007.

## Literature Cited

- Barrett, S.A., and E.W. Gifford. 1933. Miwok material culture: Indian life of the Yosemite region. Yosemite Association, Yosemite National Park, CA.
- Barton, B.J. 1997. Gardening by mail: a source book. Houghton Mifflin, Boston & New York.
- Bartow, A., editor. 2015. Native seed production manual for the Pacific Northwest. USDA-Natural Resources Conservation Service, Corvallis Plant Materials Center, Corvallis, OR.
- Brenzel, K.N., editor. 1995. Sunset western garden book. Sunset Publishing Corp., Menlo Park, CA.
- Bryan, J. 1989. Bulbs. Timber Press, Portland, OR.
- Craighead, J.J., F.C. Craighead, and R.J. Davis. 1963. A field guide to Rocky Mountain wildflowers. Houghton Mifflin, Boston, MA.
- Cooke, S.S., editor. 1997. A field guide to the common wetland plants of western Washington and northwestern Oregon. Seattle Audubon Society and Washington Native Plant Society, Seattle, WA.
- Deno, N.C. 1993. Seed germination theory and practice. Self-published and self-distributed, Professor Emeritus of Chemistry, State College, PA.
- De Hertogh, A.A., L.B. Gallitano, G.H. Pemberton, and M.E. Traer. 1993. Guidelines for the utilization of flowering bulbs as perennial (naturalized) plants in North American landscapes and gardens. Bulletin No. 37. Holland flower bulb technical services, Netherlands Flower Bulb Center (IBC), Hillegom, The Netherlands.

- De Hertogh, A.A., and M. De Nard. 1993. General chapter on spring flowering bulbs. In: De Hertogh, A.A., and M. De Nard, editors, *The physiology of flower bulbs: a comprehensive treatise on the physiology and utilization of ornamental flowering bulbous and tuberous plants*. Elsevier Science, New York, NY. p. 705-707.
- De Hertogh, A.A., C. Noone, and A. Lutman. 1990. *Camassia*. In: Geophyte, ver. 2.2. North Carolina State Univ., Dept. of Horticultural Science, Raleigh. p. 8.
- Emery, D.E. 1988. Seed propagation of native California plants. Santa Barbara Botanic Garden, Santa Barbara, CA.
- Guard, J.B. 1995. Wetland plants of Oregon and Washington. Lone Pine Publishing, Redmond, WA, Vancouver, B.C., & Edmonton, Alberta.
- Guerrant, E.O. Jr., and A. Raven. 1995. Seed germination and storability studies of 69 plant taxa native to the Willamette Valley wet prairie. The Berry Botanic Garden, Portland, OR.
- Gunther, E. 1945. Ethnobotany of western Washington, rev. 1973. Univ. of Washington Publications in Anthropology 10(1). Univ. of Washington Press, Seattle.
- Harbinger, L.J. 1964. The importance of food plants in the maintenance of Nez Perce cultural identity. Thesis, Master of Arts in Anthropology, Washington State Univ., Pullman.
- Hickman, J.C., editor. 1993. *The Jepson manual: higher plants of California*. Univ. of California Press, Berkeley.
- Hitchcock, C.L., and A. Cronquist, editors. 1973. *Flora of the Pacific Northwest: an illustrated manual*. Univ. of Washington Press, Seattle.
- International Flower Bulb Center. 2017. *Camassia leichtlinii*. Technical information service of the IFBC, Hillegom, The Netherlands. <http://us.bulb.com/836/camassia-leichtlinii.html> (accessed 17 Aug. 2017).
- Isaacson, R.T. 1993. Anderson horticultural library's source list of plants and seeds. Anderson Horticultural Library, University of Minnesota Libraries. Minnesota Landscape Arboretum.
- Kruckeberg, A.R. 1996. *Gardening with native plants of the Pacific Northwest*, 2nd edition. Univ. of Washington Press, Seattle.
- Kuhnlein, H.V., and N.J. Turner. 1991. Traditional plant foods of Canadian indigenous peoples: nutrition, botany, and use. Vol. 8 In: S.H. Katz, editor, *Food and nutrition in history and anthropology*. Gordon and Breach Science Publishers, Amsterdam, The Netherlands.
- Langaslag, J.J.J. 1989. *Camassia*. In: *Teelt en gerbruiksmogelijkheden van bijgoedgewassen*. Tweede Druk. Ministerie Landbouw Visserij en Consulentschap Algemene Dienst Bloembollenteelt, Lisse, The Netherlands. p. 76-78.
- Martin, A.C., H.S. Zim, and A.L. Nelson. 1951. *American wildlife and plants: a guide to wildlife food habits*. Dover Publications, New York, NY.
- Mayer, K.E., and W.F. Laudenslayer Jr., editors. 1988. *A guide to wildlife habitats of California*. USDA, Forest Service and California Dept. of Fish and Game, Sacramento.
- Murphey, E.V.A. 1959. *Indian uses of native plants*. Mendocino County Historical Society, Ukiah, CA.
- Pendergrass, K., M. Vaughan, and J. Williams. 2008. *Plants for pollinators in Oregon*. Plant Materials Technical Note No. 13. USDA-Natural Resources Conservation Service, Portland, OR.
- Piper, C.V. 1916. Notes on Quamasia with a description of a new species. *Proceedings of the Biological Society of Washington* 24:77.
- Pojar, J., and A. MacKinnon, editors. 1994. *Plants of the Pacific Northwest coast: Washington, Oregon, British Columbia and Alaska*. British Columbia Ministry of Forests and Lone Pine Publishing.
- Schopmeyer, technical coordinator. 1974. *Seeds of woody plants in the United States*. Agriculture Handbook No. 450. USDA, Forest Service, Washington, D.C.
- Showers, M.A., and K. Wiese, editors. 1995. *Nursery sources for California native plants*. DMG Open File Report 90-04. California Dept. of Conservation, Office of Mine Reclamation, Sacramento, CA.
- Thoms, A. 1989. The northern roots of hunter-gatherer intensification: camas and the Pacific Northwest. Ph.D. diss., Dept. of Anthropology, Washington State Univ., Pullman.
- Turner, N.J., and M.A.M. Bell. 1983. The ethnobotany of the Southern Kwakiutl Indians of British Columbia. *Econ. Bot.* 27:257-310.
- Turner, N.J., and B.S. Efrat. 1982. The ethnobotany of the Hesquiat Indians of Vancouver Island. Cultural Recovery Paper No. 2, British Columbia Provincial Museum, Queen's Printer, Victoria, BC.
- Turner, N.J., and H.V. Kuhnlein. 1983. Camas (*Camassia* spp.) and riceroot (*Fritillaria* spp.): two liliaceous "root" foods of the northwest coast Indians. *Ecology of Food and Nutrition* 13:199-219.
- Watson, R.D. 1988. Interview data on Camas growing in Moscow, Idaho. Interviewed (4-18-88) by Alson V. Thoms. Notes on file, Center for Northwest Anthropology, Washington State Univ., Pullman.

## Citation

Stevens, M., D.C. Darris, and S.M. Lambert. 2000. Plant guide for common camas (*Camassia quamash* ssp. *breviflora*). USDA-Natural Resources Conservation Service, National Plant Data Center, Greensboro, NC, and Corvallis Plant Materials Center, Corvallis, OR.

Edited: 05Dec2000 jsp; 01May2003 ahv; 31May2006 jsp; 31Aug2017 aym

For more information about this and other plants, please contact your local NRCS field office or Conservation District at <http://www.nrcs.usda.gov/> and visit the PLANTS Web site at <http://plants.usda.gov/> or the Plant Materials Program Web site: <http://plant-materials.nrcs.usda.gov>.

PLANTS is not responsible for the content or availability of other Web sites.

The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers, employees, and applicants for employment on the bases of race, color, national origin, age, disability, sex, gender identity, religion, reprisal, and where applicable, political beliefs, marital status, familial or parental status, sexual orientation, or all or part of an individual's income is derived from any public assistance program, or protected genetic information in employment or in any program or activity conducted or funded by the Department. (Not all prohibited bases will apply to all programs and/or employment activities.)

If you wish to file an employment complaint, you must contact your agency's [EEO Counselor](#) (PDF) within 45 days of the date of the alleged discriminatory act, event, or in the case of a personnel action. Additional information can be found online at [http://www.ascr.usda.gov/complaint\\_filing\\_file.html](http://www.ascr.usda.gov/complaint_filing_file.html).

If you wish to file a Civil Rights program complaint of discrimination, complete the [USDA Program Discrimination Complaint Form](#) (PDF), found online at [http://www.ascr.usda.gov/complaint\\_filing\\_cust.html](http://www.ascr.usda.gov/complaint_filing_cust.html), or at any USDA office, or call (866) 632-9992 to request the form. You may also write a letter containing all of the information requested in the form. Send your completed complaint form or letter to us by mail at U.S. Department of Agriculture, Director, Office of Adjudication, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410, by fax (202) 690-7442 or email at [program.intake@usda.gov](mailto:program.intake@usda.gov). Individuals who are deaf, hard of hearing or have speech disabilities and you wish to file either an EEO or program complaint please contact USDA through the Federal Relay Service at (800) 877-8339 or (800) 845-6136 (in Spanish).

Persons with disabilities who wish to file a program complaint, please see information above on how to contact us by mail directly or by email. If you require alternative means of communication for program information (e.g., Braille, large print, audiotope, etc.) please contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).

For any other information dealing with Supplemental Nutrition Assistance Program (SNAP) issues, persons should either contact the USDA SNAP Hotline Number at (800) 221-5689, which is also in Spanish or call the [State Information/Hotline Numbers](#). For any other information not pertaining to civil rights, please refer to the listing of the [USDA Agencies and Offices](#) for specific agency information.

*Helping People Help the Land*

**USDA IS AN EQUAL OPPORTUNITY PROVIDER, EMPLOYER, AND LENDER**